
The Forgotten Electrolyte in Dengue: Hypophosphatemia Matters

Karan Chaman Lal¹, Syeda Zainab Kazmi², Sana Soomro³, Ayla Tariq², Ayesha Ubaid Ullah²

¹Department of Medicine, Liaquat University of Medical & Health Sciences, Jamshoro, Sindh, Pakistan

²Department of Medicine, Islamic International Medical College, Rawalpindi, Pakistan.

³Department of Medicine, Jinnah Sindh Medical University, Karachi, Sindh, Pakistan.

¹Corresponding Author Email: Karan.chaman12@gmail.com

KEYWORDS:

Dengue, Hypophosphatemia, Electrolyte imbalance, Serum phosphate, Hemorrhagic fever.

Dear Editor,

Dengue is an acute viral illness from the family Flaviviridae that has caused significant morbidity and mortality worldwide, with a broad disease spectrum, ranging from asymptomatic infection to severe dengue hemorrhagic fever and dengue shock syndrome. With 2.5 billion people at risk globally, especially in endemic regions, the burden continues to rise, underscoring the need for early recognition and comprehensive monitoring as crucial parts of patient care (1).

Although various laboratory investigations have been performed to assess the severity and progression of disease, such as complete blood count (CBC), prothrombin time (PT), activated partial thromboplastin time (aPTT), and liver function test (LFT) (2), and serum electrolytes, mainly sodium and potassium, serum phosphate levels are almost entirely ignored despite their high clinical relevance. A pediatric study published in the Pakistan Journal of Medicine and Dentistry reported that 65.3% of dengue patients had decreased serum phosphate levels (3), highlighting the wide gap in critical care and the current literature, where only selected electrolytes are monitored, leaving unmonitored serum phosphate alterations unnoticed. However, adult data remains limited, highlighting the need for further prospective studies.

Hypophosphatemia is defined as a serum phosphate level below 0.80 mmol/L ($\approx < 2.5$ mg/dL). It is further classified as mild (0.60–0.79 mmol/L ≈ 1.9 –2.4 mg/dL), moderate (0.32–0.59 mmol/L ≈ 1.0 –1.8 mg/dL), and severe (< 0.32 mmol/L $\approx < 1.0$ mg/dL) (4). Decreased serum phosphate levels are commonly observed in critically ill patients and have also been reported in patients with dengue infection. However, various causes, such as reduced intestinal phosphate absorption, increased renal excretion, osteolysis, intracellular changes, and metabolic abnormalities, can also result in hypophosphatemia. Numerous deleterious clinical outcomes have been observed due to low serum phosphate levels, including more extended hospital and intensive care unit stays and higher death rates because phosphate is primarily an intracellular anion that is essential for regular cellular activity, including nucleic acid structural proteins, important cellular energy sources, red blood cells, and bones and its depletion can cause severe adverse

effects such as irregular heart rhythm, hemolysis, delirium, convulsions, prolonged mechanical ventilation, coma, peripheral neuropathy, rhabdomyolysis, skeletal muscle weakness, and respiratory insufficiency (5). A case report by Nirada Siriyakorn observed fatal rhabdomyolysis in a patient with dengue hemorrhagic fever, linking hypophosphatemia as a potential mechanism of causing such illness (6). Phosphate is a primary electrolyte in genetic and metabolic pathways and drives mineralization of skeletal tissues (7). Moreover, this electrolyte plays a vital role in maintaining body homeostasis via FGF23, vitamin D, and parathyroid hormone, influencing energy balance, mineralization, rickets prevention, and treatment of phosphate-related disorders (8). It is pertinent to note that in critical illness, hypophosphatemia correlates with worse outcomes; whether targeted correction improves outcomes in dengue remains to be studied (9). Therefore, this issue warrants urgent attention from health authorities because changes in serum phosphate levels were not random but showed a consistent pattern in dengue fever.

In conclusion, several practical approaches should be implemented, such as monitoring serum phosphate levels in conjunction with dengue phases; baseline measurements on admission, daily during the critical phase, and serial tracking in the ICU or for rhabdomyolysis risk. Critical-phase alarms should serve as reminders to perform serum phosphate testing. Serum phosphate laboratory investigations should be integrated with existing chemistry panels, guided by clinical triggers. Specifically, the serum phosphate with platelet bundle test is prompted by a fall in platelet levels, requiring a single lab test co-ordering CBC with platelets and serum phosphate, thus reducing phlebotomy and costs. During the replacement protocol, oral phosphate should be preferred for mild to moderate cases, and intravenous phosphate for severe cases, with close monitoring of electrolytes and renal function to prevent complications. Therefore, integrating diligent strategies in critical care may reduce the length of hospital stay and potentially be life-saving.

Authors Contribution:

Conceptualization, K.C.L.; Original Draft, K.C.L.; Literature Review, K.C.L., S.Z.K., S.S., A.T.; Writing - Review & Editing, K.C.L., S.Z.K., S.S., A.T., A.U.U.; Final Approval, K.C.L., S.Z.K., S.S., A.T., A.U.U.

Declaration of Interests:

The authors declare no conflict of interest.

Acknowledgments:

The authors declare no acknowledgments.

Funding: Not applicable

REFERENCES

1. Hasan S, Jamdar SF, Alalowi M, Al Ageel Al Beaiji SM. Dengue virus: A global human threat: Review of literature. *J Int Soc Prev Community Dent* [Internet]. 2016 Jan 1 [cited 2025 Sep 30];6(1):1–6. Available from: <https://pubmed.ncbi.nlm.nih.gov/27011925/>

2. Yuan K, Chen Y, Zhong M, Lin Y, Liu L. Risk and predictive factors for severe dengue infection: A systematic review and meta-analysis. PLoS One [Internet]. 2022 Apr 1 [cited 2025 Sep 30];17(4):e0267186. Available from: <https://pmc.ncbi.nlm.nih.gov/articles/PMC9012395/>
3. Altered Electrolyte Homeostasis and the Severity of Dengue in Children. Pakistan J Med Dent [Internet]. 2021 May 31 [cited 2025 Sep 30]; Available from: <https://pjmd.zu.edu.pk/volume-10-issue-2/altered-electrolyte-homeostasis-and-the-severity-of-dengue-in-children/>
4. Geerse DA, Bindels AJ, Kuiper MA, Roos AN, Spronk PE, Schultz MJ. Treatment of hypophosphatemia in the intensive care unit: A review. Crit Care [Internet]. 2010 Aug 3 [cited 2025 Oct 28];14(4):1–8. Available from: <https://ccforum.biomedcentral.com/articles/10.1186/cc9215>
5. Ramanan M, Tabah A, Affleck J, Edwards F, White KC, Attokaran A, et al. Hypophosphataemia in Critical Illness: A Narrative Review. J Clin Med [Internet]. 2024 Dec 1 [cited 2025 Sep 30];13(23). Available from: <https://pubmed.ncbi.nlm.nih.gov/39685625/>
6. Siriyakorn N, Insiripong S. FATAL RHABDOMYOLYSIS IN DENGUE HEMORRHAGIC FEVER: A CASE REPORT. Southeast Asian J Trop Med Public Health [Internet]. 2015 [cited 2025 Oct 28];46 Suppl 1:149–52. Available from: <https://pubmed.ncbi.nlm.nih.gov/26506741/>
7. Bhadada SK, Rao SD. Role of Phosphate in Biomineralization. Calcif Tissue Int [Internet]. 2021 Jan 1 [cited 2025 Sep 30];108(1):32–40. Available from: <https://pubmed.ncbi.nlm.nih.gov/32712778/>
8. Michigami T. Advances in understanding of phosphate homeostasis and related disorders. Endocr J [Internet]. 2022 [cited 2025 Sep 30];69(8):881–96. Available from: <https://pubmed.ncbi.nlm.nih.gov/35831119/>
9. Wang L, Xiao C, Chen L, Zhang X, Kou Q. Impact of hypophosphatemia on outcome of patients in intensive care unit: a retrospective cohort study. BMC Anesthesiol [Internet]. 2019 May 24 [cited 2025 Sep 30];19(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/31122196/>